

On the Spectrum of the Orr-Sommerfeld Equation on the semi-axis

By J. LAHMANN, M. PLUM

Abstract. The Orr-Sommerfeld equation is a spectral problem which is known to play an important role in hydrodynamic stability. For an appropriate operator theoretical realization of the equation, we will determine the essential spectrum, and calculate an enclosure of the set of all eigenvalues by elementary analytical means.

1. Introduction

The *Orr-Sommerfeld* equation is an non-selfadjoint general eigenvalue problem of the form

$$(1.1) \quad (-D^2 + a^2)^2 u + iaR[V \cdot (-D^2 + a^2)u + V'' \cdot u] = \lambda(-D^2 + a^2)u \quad \text{on } I,$$

subject to Dirichlet boundary conditions for u on the real interval I . Here, $D := d/dx$, i is the imaginary unit, and $R > 0$ is the Reynolds number of an underlying fluid which moves in a stationary flow (perpendicular to I) with given real-valued flow profile $V \in C^2(I)$. This flow is perturbed by a single-mode perturbation with wave number $a > 0$, and the physical question of stability or instability of the underlying flow in response to this perturbation arises. This question is closely related to the spectrum of (an appropriate operator-theoretical realization of) the Orr-Sommerfeld equation (1.1). Essentially the flow is unstable (with respect to the wave number a) if (1.1) has an eigenvalue with negative real part.

In this article, we will exclusively be concerned with the case $I = [0, \infty)$ corresponding to the half-plane flow along a wall (an overview of results for this case can be found in [3]). So the boundary conditions to be added to (1.1) read

$$(1.2) \quad u(0) = u'(0) = u(\infty) = u'(\infty) = 0.$$

A flow profile V which is of particular interest in this case (but not under exclusive concern here) is the *Blasius* profile defined by $V := f'$, where f is the solution of the

nonlinear boundary value problem (Blasius equation, a special case of the Falkner–Skan equation)

$$(1.3) \quad 2f''' + ff'' = 0 \quad \text{on } [0, \infty), \quad f(0) = f'(0) = 0, \quad \lim_{x \rightarrow \infty} f'(x) = 1,$$

which can be shown to exist and to be unique, and moreover, to provide (for $V = f'$)

$$(1.4) \quad \begin{aligned} &V \rightarrow 1, \quad V' \rightarrow 0, \quad V'' \rightarrow 0 \text{ (exponentially) as } x \rightarrow \infty, \\ &V > 0, \quad V' > 0, \quad V'' < 0 \text{ on } (0, \infty). \end{aligned}$$




Figure 1: Blasius profile together with its first two derivatives

We choose the following operator theoretical setting for problem (1.1), (1.2), which we believe is the most natural and simple one:

Let X and Y denote the complex Banach spaces $H^2(0, \infty) \cap H_0^1(0, \infty)$ and $L^2(0, \infty)$, respectively, and let $D(A) := H^4(0, \infty) \cap H_0^2(0, \infty)$. The operators $A : D(A) \subset X \rightarrow Y$ and $B : X \rightarrow Y$ are defined by Au and Bu denoting the left-hand and the right-hand side (without λ) of equation (1.1), respectively. It can be shown that A is closed (see [1, Theorem IX.9.5], plus the remark that the norm in X is stronger than the one in $L^2(0, \infty)$), and that B is one-to-one and onto, and bounded with bounded inverse. Our formulation of (1.1), (1.2) now reads

$$(1.5) \quad Au = \lambda Bu.$$

So we choose a *direct* operator theoretical realization of the Orr–Sommerfeld problem, rather than a formulation via an auxiliary operator constructed by Riesz’s representation lemma, as done e.g. in [2].

Our goal in this article is twofold: In Section 2, we will determine the essential spectrum of problem (1.5) exactly; the main tool is the well-known theorem on preservation of the essential spectrum under relatively compact perturbations. In Section 3, an enclosure for the set of all eigenvalues of problem (1.5) is calculated by elementary

analytical means. Here, exploiting sign restrictions on the profile V and its derivatives (such as (1.4)), we obtain results which extend enclosures known in the literature (see [2], [5]), even if those have been obtained by more involved methods. In [4] corresponding results have been obtained for the case of a compact interval I . For further results on the Orr–Sommerfeld equation on a compact interval see [6] and the literature cited there.

2. The essential spectrum

Since several definitions of essential spectra are around in the literature (see [1, Chapter IX] for an overview), we start with the definition which we are using.

With X, Y denoting two complex Banach spaces, and $A : D(A) \subset X \rightarrow Y$ a closed linear and $B : X \rightarrow Y$ a bounded linear operator, we call the set

$$\sigma_{\text{ess}}(A, B) := \{\lambda \in \mathbb{C} : A - \lambda B \text{ is not a Fredholm operator of index } 0\}$$

the *essential spectrum* of the problem $Au = \lambda Bu$ or of the pencil (A, B) . Here, a closed linear operator $T : D(T) \subset X \rightarrow Y$ is called a *Fredholm operator* if its range is closed and has finite codimension $d(T)$ in Y , and if its nullspace has finite dimension $n(T)$; it has index 0 if $d(T) = n(T)$. Furthermore,

$$\varrho(A, B) := \{\lambda \in \mathbb{C} : A - \lambda B \text{ is one-to-one and onto}\}$$

denotes the *resolvent set* of the problem $Au = \lambda Bu$ (or of the pencil (A, B)), and $\sigma(A, B) := \mathbb{C} \setminus \varrho(A, B)$ its (total) *spectrum*. It is quite obvious that $\sigma_{\text{ess}}(A, B) \subset \sigma(A, B)$ and that $\sigma(A, B) \setminus \sigma_{\text{ess}}(A, B)$ consists only of the eigenvalues of finite geometric multiplicity.

We return to the specific choice for X, Y, A, B introduced in Section 1, i.e., describing the Orr–Sommerfeld problem. Our main result of the present section is

Theorem 2.1. *Suppose that the flow profile V satisfies $V \rightarrow 1, V'' \rightarrow 0$ as $x \rightarrow \infty$ (compare (1.4)). With R and a denoting the parameters in (1.1), the essential spectrum of (1.5) is given by*

$$(2.1) \quad \sigma_{\text{ess}}(A, B) = \{\mu + a^2 + iaR : \mu \in [0, \infty)\}.$$

Proof. We introduce the auxiliary operators

$$A_0 : D(A) \rightarrow Y, A_0 u := (-D^2 + a^2)^2 u + iaR(-D^2 + a^2)u,$$

$$K : D(A) \rightarrow Y, Ku := iaR[(V - 1) \cdot (-D^2 + a^2)u + V'' \cdot u]$$

so that $A = A_0 + K$. Since $V - 1$ and V'' tend to zero and K is of lower order than A_0 , it follows from [1, Theorem IX.8.2] that K is relatively compact to A_0 , i.e., for each $\|\cdot\|_X$ -bounded sequence (u_n) in $D(A)$ such that $(A_0 u_n)$ is bounded in Y , (Ku_n) contains a convergent subsequence. Strictly speaking, the cited Theorem in [1]

provides the result with X and $D(A)$ replaced by $L^2(0, \infty)$ and $H^4(0, \infty)$, respectively, but from this it is easy to obtain the same statement with X and $D(A)$.

Since $B : X \rightarrow Y$ is one-to-one, onto, and bounded with bounded inverse, the above result implies that $B^{-1}K : D(A) \rightarrow X$ is relatively compact to $B^{-1}A_0 : D(A) \rightarrow X$. Now the invariance of the essential spectrum under relatively compact perturbations (see e.g. [1, Theorem IX.2.1]) provides

$$(2.2) \quad \sigma_{\text{ess}}(B^{-1}A_0 + B^{-1}K, id_X) = \sigma_{\text{ess}}(B^{-1}A_0, id_X).$$

The cited theorem in [1] is formulated for densely defined operators, but the proof clearly does not use this assumption. (This density is a general assumption in [1, Chapter IX] to be able to work with adjoint operators, which however do not occur in Theorem IX.2.1 or in its proof.) Since B and B^{-1} are bounded, so that $A - \lambda B$ is a Fredholm operator of index 0 if and only if $B^{-1}A - \lambda id_X$ is, we obtain furthermore $\sigma_{\text{ess}}(A, B) = \sigma_{\text{ess}}(B^{-1}A, id_X)$, so that (2.2) and the identity $A_0 + K = A$ provide

$$(2.3) \quad \sigma_{\text{ess}}(A, B) = \sigma_{\text{ess}}(B^{-1}A_0, id_X).$$

It remains to be shown that $\sigma_{\text{ess}}(B^{-1}A_0, id_X)$ equals the right-hand side of (2.1) which we call M from now on. This is done in two steps: We prove

- a) $\mathbb{C} \setminus M \subset \varrho(B^{-1}A_0, id_X)$ (implying $M \supset \sigma(B^{-1}A_0, id_X) \supset \sigma_{\text{ess}}(B^{-1}A_0, id_X)$).
- b) $M \subset \sigma_{\text{ess}}(B^{-1}A_0, id_X)$.

First we observe that, with $E(x) := e^{-ax}$,

$$(2.4) \quad B^{-1}A_0 u = -u'' + (a^2 + iaR)u + u''(0) \cdot E \quad \text{for } u \in D(A),$$

since the expression on the right-hand side is in X (for $u \in D(A)$) and B applied to it equals $A_0 u$.

ad a) Let $\lambda \in \mathbb{C} \setminus M$, so that $\lambda = \mu + a^2 + iaR$ with $\mu \in \mathbb{C} \setminus [0, \infty)$. Thus, $\mu \in \varrho(C, id_Y)$ where $C : D(C) \subset Y \rightarrow Y$ is given by $D(C) := H^2(0, \infty) \cap H_0^1(0, \infty)$, $Cu := -u''$.

To prove that $B^{-1}A_0 - \lambda id_X : D(A) \rightarrow X$ is onto, let $r \in X$ and define $v := (C - \mu)^{-1}r \in D(C) = X$, $w := (C - \mu)^{-1}E \in D(C) = X$. It is easy to calculate w in closed form and to show that $w'(0) \neq 0$. Therefore, $u := v - \frac{v'(0)}{w'(0)}w \in X$ solves the equation

$$(2.5) \quad -u'' - \mu u + \frac{v'(0)}{w'(0)}E = r$$

and moreover, $u(0) = u'(0) = 0$. Since $u, E, r \in H^2(0, \infty)$, (2.5) implies $u'' \in H^2(0, \infty)$. Therefore $u \in H^4(0, \infty)$. Altogether, $u \in D(A)$. Since $u(0) = r(0) = 0$ and $E(0) = 1$, (2.5) shows that $u''(0) = v'(0)/w'(0)$, and (2.5), (2.4) then yield $(B^{-1}A_0 - \lambda)u = r$.

To prove that $B^{-1}A_0 - \lambda id_X$ is one-to-one, let $u \in D(A)$ satisfy $(B^{-1}A_0 - \lambda)u = 0$, i.e.,

$$-u'' - \mu u + u''(0)E = 0.$$

Thus, $u \in D(C)$ and $(C - \mu)u = -u''(0)E$, i.e., $u = -u''(0)w$. Since $u'(0) = 0$, $w'(0) \neq 0$, this implies $u''(0) = 0$ and therefore $u = 0$.

ad b) Let $\lambda \in M$, so that $\lambda = \mu + a^2 + iaR$ with $\mu \in [0, \infty)$. Let $\gamma \in \mathbb{C}$ and $r := \gamma(1 - E)E \in X$, and suppose that, for some $u \in D(A)$, $(B^{-1}A_0 - \lambda)u = r$, so that (2.4) yields

$$(2.6) \quad -u'' - \mu u + u''(0)E = \gamma(1 - E)E.$$

We will prove that (2.6), together with the fact $u \in D(A)$, implies $\gamma = 0$ and $u = 0$, which shows that $B^{-1}A_0 - \lambda \text{id}_X$ is one-to-one but not onto. Consequently, it is no Fredholm operator of index zero, which implies $\lambda \in \sigma_{\text{ess}}(B^{-1}A_0, \text{id}_X)$. The general solution of (2.6) satisfies

$$(2.7) \quad u = c_1\varphi_1 + c_2\varphi_2 + \frac{u''(0) - \gamma}{a^2 + \mu}E + \frac{\gamma}{4a^2 + \mu}E^2$$

with $c_1, c_2 \in \mathbb{C}$, where $\varphi_1(x) = \cos(\sqrt{\mu}x)$ and $\varphi_2(x) = \sin(\sqrt{\mu}x)$ if $\mu > 0$, $\varphi_1(x) = 1$ and $\varphi_2(x) = x$ if $\mu = 0$. The condition $u \in L^2(0, \infty)$ requires $c_1 = c_2 = 0$, and $u(0) = u'(0) = 0$ then implies

$$\frac{\gamma - u''(0)}{a^2 + \mu} = \frac{\gamma}{4a^2 + \mu}, \quad a \frac{\gamma - u''(0)}{a^2 + \mu} = 2a \frac{\gamma}{4a^2 + \mu}$$

which indeed yields $\gamma = 0$, $u''(0) = 0$, so that (2.7) provides $u \equiv 0$. $\square$

It is easy to generalize Theorem 2.1 for flow profiles which satisfy $V \rightarrow c \in \mathbb{R}$ instead of $V \rightarrow 1$.

Corollary 2.2. *Suppose that the flow profile V satisfies $V \rightarrow c \in \mathbb{R}$, $V'' \rightarrow 0$ as $x \rightarrow \infty$. Then the essential spectrum of (1.5) is given by*

$$(2.8) \quad \sigma_{\text{ess}}(A, B) = \{\mu + a^2 + iaR : \mu \in [0, \infty)\}.$$

3. An enclosure for the set of all eigenvalues

In this final section, we will calculate a set enclosing all eigenvalues of problem (1.5). Since this set will also contain the straight line which was identified as $\sigma_{\text{ess}}(A, B)$ in Theorem 2.1, it therefore encloses the *total* spectrum $\sigma(A, B)$.

We wish to put emphasize on the simplicity of the methods we use, which nevertheless provide more accurate enclosures than those known in the literature [2], [5], if sign restrictions such as (1.4) are exploited.

Let real constants $V_{\min}, V_{\max}, |V'|_{\max}, V''_{\min}, V''_{\max}$ be given such that

$$(3.1) \quad V_{\min} \leq V(x) \leq V_{\max}, |V'(x)| \leq |V'|_{\max}, V''_{\min} \leq V''(x) \leq V''_{\max} \quad \text{for } x \in [0, \infty)$$

for the flow profile V . Let $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ denote the usual inner product and norm in $L^2(0, \infty)$.

For any eigenpair $(u, \lambda) \in D(A) \times \mathbb{C}$ of problem (1.5) we obtain (note that B is positive definite)

$$(3.2) \quad \lambda = \frac{\langle Au, u \rangle}{\langle Bu, u \rangle}$$

which is the basis of our enclosures. Partial integration and the condition $u \in D(A) = H^4(0, \infty) \cap H_0^2(0, \infty)$ provide

$$\begin{aligned} \langle Au, u \rangle &= \langle (-D^2 + a^2)^2 u, u \rangle + iaR \langle V \cdot (-D^2 + a^2)u + V'' \cdot u, u \rangle \\ &= \|(-D^2 + a^2)u\|^2 + iaR[\langle Vu', u' \rangle - \langle V'u, u' \rangle + a^2 \langle Vu, u \rangle] \end{aligned}$$

so that (3.2) yields

$$(3.3) \quad \lambda = \beta_1 + iaR(\beta_2 - \beta_3)$$

where

$$(3.4) \quad \beta_1 := \frac{\|Bu\|^2}{\langle Bu, u \rangle} \geq \frac{\|Bu\|}{\|u\|} \geq a^2,$$

$$\begin{aligned} (3.5) \quad \beta_2 &:= \frac{\langle Vu', u' \rangle + a^2 \langle Vu, u \rangle}{\langle Bu, u \rangle} = V(\xi) \frac{\langle u', u' \rangle + a^2 \langle u, u \rangle}{\langle Bu, u \rangle} \\ &= V(\xi) \in [V_{\min}, V_{\max}] \quad (\text{for some } \xi \in (0, \infty)), \end{aligned}$$

$$(3.6) \quad \beta_3 := \frac{\langle V'u, u' \rangle}{\langle Bu, u \rangle}, \quad |\beta_3| \leq |V'|_{\max} \frac{\|u\| \|u'\|}{\|u'\|^2 + a^2 \|u\|^2} \leq \frac{|V'|_{\max}}{2a}.$$

From (3.3) to (3.6) we obtain the following eigenvalue enclosure result, where sums and products of sets are to be understood in the canonical sense.

Theorem 3.1. *All eigenvalues of problem (1.5) are contained in the set*

$$[a^2, \infty) + iaR[V_{\min}, V_{\max}] + \frac{R}{2}|V'|_{\max} \cdot \Delta,$$

with Δ denoting the closed unit disc in $\mathbb{C}$.

Of course, separate bounds for the real and imaginary parts of the eigenvalues can easily be extracted (with loss of information!) from Theorem 3.1:

Corollary 3.2. *For each eigenvalue λ of problem (1.5),*

$$\begin{aligned} a^2 - \frac{R}{2}|V'|_{\max} &\leq \operatorname{Re} \lambda \\ aRV_{\min} - \frac{R}{2}|V'|_{\max} &\leq \operatorname{Im} \lambda \leq aRV_{\max} + \frac{R}{2}|V'|_{\max}. \end{aligned}$$

Figures 2 and 3 illustrate the results of Theorem 3.1, Corollary 3.2 and Theorem 3.3. Concretely, we have chosen V to be the Blasius profile and $a = 0.179, R = 580$ here.




Figure 2: Illustration of Theorem 3.1 and Corollary 3.2

The bounds given in Corollary 3.2 have been obtained in [2] already, by more involved analytical means. These results can be substantially improved by the following simple calculation, if sign restrictions such as (1.4) are used:

$$\begin{aligned} \operatorname{Re} \langle V' u, u' \rangle &= \frac{1}{2} (\langle V' u, u' \rangle + \overline{\langle V' u, u' \rangle}) = \frac{1}{2} \int_0^\infty V' (|u|^2)' dx \\ &= -\frac{1}{2} \int_0^\infty V'' |u|^2 dx = -\frac{1}{2} V''(\xi) \langle u, u \rangle \quad (\text{for some } \xi \in (0, \infty)) \end{aligned}$$

so that β_3 defined in (3.6) satisfies

$$(3.7) \quad \operatorname{Re} \beta_3 = -\frac{1}{2} V''(\xi) \frac{\langle u, u \rangle}{\langle Bu, u \rangle} \in -\frac{1}{2} [V''_{\min}, V''_{\max}] \cdot [0, \frac{1}{a^2}]$$

which possibly restricts the enclosure set for β_3 used in Theorem 3.1 (which was obtained from (3.6)). Instead of formulating a corresponding improved theorem for the most general situation, we concentrate on the case where $V'' \leq 0$, which is true e.g. for the Blasius profile (see (1.4)). Then (3.7) yields $\operatorname{Re} \beta_3 \geq 0$, which together with (3.3) to (3.6) provides the following improvement of Theorem 3.1:

Theorem 3.3. *If $V'' \leq 0$, all eigenvalues of problem (1.5) are contained in the set*

$$[a^2, \infty) + iaR[V_{\min}, V_{\max}] + \frac{R}{2}|V'|_{\max} \cdot \Delta^-,$$

where $\Delta^- := \{z \in \Delta : \operatorname{Im} z \leq 0\}$.

As in Corollary 3.2, we can extract separate bounds for real and imaginary parts of the eigenvalues, which provides here the improved bound

$$aRV_{\min} - \frac{R}{2}|V'|_{\max} \leq \operatorname{Im} \lambda \leq aRV_{\max}.$$

In particular all eigenvalues lie "below" the essential spectrum, i.e. have imaginary parts less or equal aR .




Figure 3: Illustration of Theorem 3.3

Acknowledgements

The authors are grateful to Andrei Shkalikov for helpful comments and suggestions.

References

- [1] D. E. Edmunds, W. D. Evans. *Spectral Theory and Differential Operators*. Oxford University Press, New York, 1987
- [2] T. M. Fischer. *Ein mathematisch-physikalisches Modell zur Beschreibung transitioneller Grenzschichtströmungen*. DLR-Forschungsbericht 95-06, 1995
- [3] I. Herron. *The Orr-Sommerfeld equation on infinite intervals*. SIAM-Rev. 29 (1987) 597–620
- [4] D. D. Joseph. *Eigenvalue bounds for the Orr-Sommerfeld equation*. J.-Fluid-Mech. 33 (1968) 617–621
- [5] M. Miklavcic. *Stability of mean flows over an infinite flat plate*. Arch.-Rational-Mech.-Anal. 80 (1982) 57–69
- [6] A. A. Shkalikov, C. Tretter. *Spectral Analysis for linear Pencils $N - \lambda P$ of Ordinary Differential Operators*. Math. Nachr. 179 (1996) 275–305